



N2010

Reactive Nitrogen Management for Sustainable Development - Science, Technology and Policy

5th International Nitrogen Conference 2010

3rd - 7th December, New Delhi, India


Best management practices to improve fertilizer N use efficiency in Argentinean agriculture

Fernando O. Garcia and Fernando Salvagiotti

IPNI Southern Cone and EEA INTA Oliveros

[fgarcia @ipni.net](mailto:fgarcia@ipni.net) - fsalvagiotti@correo.inta.gov.ar






Grain production in Argentina

- Approximately 90% of grain crops harvested area is located in the Pampas-Chaco region, in the east-central plains of Argentina.
- Soils of the Pampas are deficient in nitrogen (N), phosphorus (P), and sulfur (S).
- 65% of the cropping area is under no-tillage
- Approximately 50% of the total cropped land is under annual leasing conditions




Field crops at Argentina Production from 1990 to 2009



Elaborated from information of MinAgri

N fertilizer consumption in Argentina 1993-2009




Source: MinAgri and Fertilizar AC



N Balance by Crop

Argentina 2009/10



Fate of N fertilizer applied to wheat and maize crops in the Pampas region of Argentina




Sink	Range	References
Plant	35 to 80%	<i>Melaj et al. 2003; Portela et al. 2006; Rillo and Richmond 2006; Rimski-Korsakov et al. 2008</i>
Organic matter	7 to 29%	<i>Sainz Rozas et al. 2004; Portela et al. 2006 ; Rimski-Korsakov et al. 2008;</i>
Volatilization	1.1 to 30%	<i>Videla et al., 1996; Garcia et al. 1999; Sainz Rozas et al. 2004; Rimski-Korsakov et al. 2007a</i>
Denitrification	0.13 to 6.9%	<i>Palma et al. 1997; Picone et al. 1997; Sainz Rosas et al. 2001; Ciampitti et al. 2008</i>
Leaching	<0.01 to 23%	<i>Sainz Rozas, et al. 2004; Portela et al. 2006 ; Aparicio et al. 2008</i>

Adapted from Lavado et al. (2007)

Soybean inoculation

A. Perticari - INTA Inocular
1994-2004



*Biological N fixation provides up to 75%
of the N accumulated by the crop*

Right N rate for wheat and maize crops

Soil testing for nitrate-N at planting (0-60 cm)

Partial N budgets

N mineralization indices (Nmin, POM)

Crop simulation models


Sap nitrate concentration in stems

*Crop canopy sensors (Minolta SPAD 502 or
remote sensors such as Greenseeker)*


Critical levels of available N at planting (NO_3^- -N, 0-60 cm depth) for wheat and maize in different areas of the Pampas with different expected yields

Area	Critical level		Expected yield + fertilizer)	Reference
	(NO_3^- -N, 0-60 cm	kg ha^{-1}		
Wheat				
SE Buenos Aires	125	3500	González Montaner et al., 1991	
SE Buenos Aires	175	5000-5500	González Montaner et al., 2003	
Central and South Santa Fe	92	3500-4000	Salvagiotti et al., 2004b	
Southern Santa Fe and Córdoba	100-150	3200-4400	García et al., 2006	
Maize				
Northern Buenos Aires	150	9000	Ruiz et al., 2001	
Northern Buenos Aires	150-170	10000	Alvarez et al., 2003	
Central and South Santa Fe	135	< 9500	Salvagiotti et al., 2004c	
	162	> 9500		
Southern Santa Fe and Córdoba	150-200	10000-11000	García et al., 2006	

Available N at planting and maize grain yield




Right time: N losses by leaching and denitrification for two different application times in wheat



Barbieri et al., 2008


Right time: Grain yield under two different application times in wheat



* denotes significant differences between treatments

Barbieri et al., 2008

Right source and place: Maize yield under different N sources and placement at central Santa Fe



Fontanetto, 2004



*Current work looking at Enhanced-Efficiency Nitrogen Sources:
Urease and nitrification inhibitors*


Urease Inhibitors in Maize

Fontanetto, Bianchini et al., 2007/08

Treatment	NH ₃ -N Losses	Grain yield	NUE
	%	kg/ha	kg grain/kg N
Check	-	7334	-
Urea 70N	10	8381	15
Urea 70N + NBTPT	4	9166	26
Urea 140N	25	9623	16
Urea 140N + NBTPT	6	10368	22

Wheat: N use efficiency without and with P application

Compiled from Senigagliesi et al. (1987) and other authors (1998-2007)





Nitrogen use efficiency (NUE), N Recovery efficiency (RE) and N Internal efficiency (IE) of wheat crops fertilized only with N (N100) and N + S (N100+ S20)

Variable	Units	N100	N100 +S20
NUE	kg grain per kg applied N	8.4	10.7
RE	kg N uptake per kg applied N	0.35	0.47
IE	kg grain per kg N uptake	22.7	22.5

Salvagiotti et al., 2009

Hairy vetch as winter crop supplying N for maize (J. Romagnoli. 2007/08)



$5000 \text{ kg DM} \rightarrow 130 \text{ kg/ha N}$

Final considerations

- ✓ Right N management for increasing the overall production of grain crops and cropping system effectiveness and efficiency, and simultaneously avoiding negative effects on the environment
- ✓ Tools for deciding the right rate, source, time, and placement of fertilizer N have been developed
- ✓ Future research should be oriented at exploring causes that may increase NUE:
 - ✓ grain yield gaps and constraints for grain production in different cropping systems
 - ✓ Nutrient recommendations should be based in more mechanistic approaches
- ✓ It is also important to look for the impact of N management on the whole system effectiveness not only in the short- but also in the long-term by watching the direct effects of nutrient addition, and also the indirect effects through increasing organic matter in soils

Thank you!!

fgarcia@ipni.net
www.ipni.net/lasc

